

R.M. Surmanidze

email: ruslan.surmanidze@tsu.ge

Mathematics department, Ivane Javakhishvili Tbilisi State University

For the class of isotropy irreducible homogeneous Riemann Manturov-Wolf spaces $M = \mathfrak{g}/\mathfrak{h}$, where the subgroup \mathfrak{h} is of the type simple Lie algebra D_n , $n \geq 2$ [1], [2], we constructing \mathfrak{g} -invariant tensors of valence 2, 3, 4 [5].

References

- [1]. O. V. Manturov, Homogeneous Riemannian spaces with an irreducible rotation group. (Russian) Trudy Sem. Vektor. Tenzor. Anal. 13 (1966), 68-145.
- [2]. J. Wolf, The geometry and structure of isotropy irreducible homogeneous spaces. Acta Math., 120, No. 1-2, 1968, pp 59-148.
- [3]. O. V. Manturov, Homogeneous spaces and invariant tensors. *Journal of Soviet Mathematics*, September, 1988, Vol. 42, **Issue 5**, pp 1944–1973.
- [4]. O. V. Manturov, Elementy Tenzornogo ischislenia. M. "Proscschenie"1991.
- [5]. R. M. Surmanidze, Tensor invariants and homogeneous Riemann spaces, *Journal of Mathematical Sciences*, November, 2013, Vol. 195, Issue 2, pp 245-257.